

CORRECTED
VERSION*

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44	A1	(11) International Publication Number: WO 99/07368 (43) International Publication Date: 18 February 1999 (18.02.99)
(21) International Application Number: PCT/US98/16262 (22) International Filing Date: 5 August 1998 (05.08.98) (30) Priority Data: 08/906,466 5 August 1997 (05.08.97) US (71)(72) Applicant and Inventor: GRASSETTI, Davide, R. [US/US]; 19810 Peppermint Falls Road, Jamestown, CA 95327 (US). (74) Agent: BROTMAN, Harris, F.; The Brotman Group, Suite 301, 7911 Herschel Avenue, La Jolla, CA 92037 (US).		(81) Designated States: AU, CA, CN, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ANTIMUTAGENIC AGENTS		
(57) Abstract A method of treating a human subject to inhibit mutagenic effects of carcinogens by administering to the subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid to inhibit activity of poly(ADP-ribose) polymerase in said subject to thereby inhibit said effects. Also provided is a method of inhibiting occurrence of mutations and of cancers derived from mutations, which method comprises administration to a subject of a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid.		

*(Referred to in PCT Gazette No. 19/1999, Section II)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ANTIMUTAGENIC AGENTS

This application is a continuation-in-part of U.S. Patent Application 08/906,466 filed August 5, 1997, which derives priority of U.S. Provisional Application 60/026,118 filed September 16, 1996.

BACKGROUND OF THE INVENTION

Field of the Invention

10 This invention relates to a method for inhibiting the mutagenic effects of carcinogens, and in particular, a method for administering to a human the compound 6,6' dithiodinicotinic acid to inhibit the activity of the enzyme poly(ADP-ribose) polymerase to inhibit the occurrence of mutation-caused cancer.

15 Background of the Invention

Mutations occur in cellular DNA under the effect of ultraviolet light, infrared light, x-rays, ionizing radiation, and chemicals. It is generally accepted that most mutations lead to cancer.

20 The enzyme poly(ADP-ribose) polymerase (ADPRP) has been suspected to play a regulatory role in many cellular activities, including DNA repair, differentiation, and malignant transformation. Studies of inhibitors of ADPRP have provided evidence of the importance of ADPRP in those cellular activities (M.J. Suto et al., 1991, *Drugs of the Future*, 16:723-739). Many analogs of nicotinamide, a product of ADPRP's action on
25 NAD, have been studied in the hope of finding a compound that could modulate the activity of ADPRP. It would be desirable to find a selective inhibitor of ADPRP which could inhibit the malignant transformation of human cells (*Ibid*).

SUMMARY OF THE INVENTION

30

It has been discovered that 6,6'-dithiodinicotinic acid has an antimutagenic effect. The method of the invention is based on the finding that 6,6'-dithiodinicotinic acid is a useful agent for the prevention of mutations, most of which lead to cancer.

The method of the invention comprises treating a human subject to inhibit the mutagenic effects of carcinogens. The method involves administering to the subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid to inhibit the activity of poly(ADP-ribose) polymerase to thereby inhibit the mutagenic effects, and in particular, their carcinogenic effects. In another aspect, the invention involves a method of inhibiting the occurrence of mutations and of cancers derived from mutations, and involves administering to a subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid.

10

DETAILED DESCRIPTION OF THE INVENTION

General Description and Definitions

The practice of the present invention will employ, unless otherwise indicated, conventional molecular and cell biology, cell culture, biochemistry, and organic and medicinal chemical synthesis within the skill of the art. Such techniques are explained fully in the literature. See Cancer Chemotherapy: Principles and Practice, ed. B.A. Chabner, J.M. Collins, Phil., Lippincott Publ., 1990; Cancer: Principles and Practice of Oncology; ed. de Vita, Jr., V.T., Hellman, S., and Rosenberg, S.A., Lippincott Co., Philadelphia, 1993; The Chemotherapy Source Book, ed. Perry, M.C., Williams and Wilkins Publ., Baltimore, 1991). Silverman, Richard B., The Organic Chemistry of Drug Design and Drug Action, Academic Press, Inc. NY (1992); Smith, Michael B., Organic Synthesis, McGraw Hill, Inc., NY, (1994)).

Methods are well known in the art for determining therapeutically effective amounts of the compounds used in the method of the invention. Such methods involve analysis of the pharmaceutical/pharmacokinetic parameters in anti-cancer or antitumor therapy, i.e for inhibiting the growth of cancerous tumors (Wedge. S.R., Porteus, J.K., Newlands, E.S., Cancer Chemother. Pharmacol. (1997) 40:266-272; Legha, S.S., Seminar in Oncology, (1997) 24:S4-24-31; Motzer, R.J., Vogelzang, N.J., Chemotherapy for Renal Cell Carcinoma. In: Raghaven, D., Scher, H.I., Leibel, S.A., et al: eds. Principles and Practice of Genitourinary Oncology, Lippincott-Raven Publ., Philadelphia, pp. 885-96, 1997; Bloom, H.J., Medroxyprogesterone acetate (Provera) in the treatment of metastatic renal cancer, Br. J. Cancer (1971) 25:250-65)

The present invention shows that in the presence of 6,6'-dithiodinicotinic acid the mutagenic effect of carcinogenic chemicals, exemplified by benzo-[a]-pyrene, 2-aflatoxin, and 2-aminofluorene, is greatly diminished.

5

6,6'-dithiodinicotinic acid (carboxypyridine disulfide, CPDS) (D.R. Grassetti, 1986, *Drugs of the Future*, 11(7):559-61) is known to prevent metastases in mice (Grassetti, D.R., 1970, *Nature* 228:282).

10 Examples 1, 2, and 3 report experiments in which the antimutagenic effects of 6,6'-dithiodinicotinic acid were observed. These experiments were performed with a mutant strain of *Salmonella Typhimurium* bacteria T-100. When the cells were exposed to varying amounts of benzo-[a]-pyrene, 2-aflatoxin, or 2-aminofluorene, a number of colonies (mutant) were formed. When 6,6'-dithiodinicotinic acid was present in the
15 culture prior to the addition of carcinogen, the number of mutants decreased by about 70%.

As shown in Example 4 below, a method of the present invention is based on the finding that 6,6'-dithiodinicotinic acid inhibited the activity of poly(ADP-ribose) polymerase
20 (ADPRP). Inhibition of ADPRP is useful for blocking the mutagenic effects of carcinogens (Suto and Suto, (1991), *Drugs of the Future*, 16(8):723-739). It is understood that 6,6'-dithiodinicotinic acid includes derivatives thereof, including, but not limited to alkaline metal salts and alkyl esters.

25 The method of the present invention includes administering a pharmaceutical composition comprising an effective amount of 6,6'-dithiodinicotinic acid in pure form or as a pharmaceutically acceptable crude concentrate in association with a pharmaceutical carrier or diluent. Such compositions conveniently contain less than 1% by weight, and preferably about 0.2% by weight, of 6,6'-dithiodinicotinic acid, and involve administration
30 of at least about 0.1 mg/kilo of body weight of 6,6'-dithiodinicotinic acid. The compositions may be prepared by conventional techniques to be in conventional forms, for example, capsules, tablets, suppositories, dispersible powders, syrups, elixirs, suspensions or solutions for enteral or parenteral administration. Suitable pharmaceutical

diluents or carriers include, for example, water, alcohols, natural or hardened oils and waxes, calcium and sodium carbonates, calcium phosphate, kaolin, talc and lactose as well as suitable preserving agents, such as ethyl-p-hydroxybenzoate, suspending agents such as methyl cellulose, tragacanth and sodium alginate, wetting agents such as lecithin, polyoxyethylene stearate and polyoxyethylene sorbitan mono-oleate, granulating and disintegrating agents such as starch and alginic acid, binding agents such as starch, gelatin and acacia and lubricating agents such as magnesium stearate, stearic acid and talc, in order to provide an elegant and palatable pharmaceutical preparation. Compositions in tablet form may be coated by conventional techniques to delay disintegration of the tablet and absorption of the active ingredient in the gastrointestinal tract and thereby provide sustained action over a long period. Other compounds and methods known in the art for delaying disintegration or for timed-delayed or time-measured delivery of the active ingredients also find use in formulating the active ingredients for use in the methods of the invention. For example, 6,6'-dithiodinicotinic acid may also be combined with liposomes or other delayed-release carrier means to protect the compounds from degradation until they reach their targets and/or facilitate movement of the compounds across tissue barriers.

The preferred compositions from the standpoint of ease of administration are solid compositions, particularly solid-filled gelatin capsules or tablets.

The carcinogens tested here are representative of the carcinogens which form a major part of air pollutants caused by tobacco smoke, the exhaust fumes of internal combustion engines, among other sources.

It has been observed that 6,6'-dithiodinicotinic acid is not a toxic compound. It can be taken in doses of at least approximately 0.1 gram per day for indefinite periods. This fills the requirements for a non-toxic, low-cost chemopreventive agent, which could be administered orally.

LEGEND TO TABLES 1, 2, 3

The effect of 6,6'-dithiodinicotinic acid (0.3 mg/plate, in 0.2 M sodium bicarbonate) added to each plate in the Ames test, which is a reverse mutation assay (Ames, B.N. et al, 1975, Mutation Research 31:347) with Salmonella Typhimurium bacteria T-100 was
 5 studied. The 6,6'-dithiodinicotinic acid was added to each plate one hour before the addition of the carcinogen. The results reported in the tables represent the average values (number of colonies per plate) of 3 individual determinations.

10 **Example 1. Effects of 6,6'-dithiodinicotinic acid on mutagenesis by benzo-[a]-pyrene**

Number of Colonies Per Plate

Benzo-[a]-pyrene (μ g/plate)	No Inhibitor	With Inhibitor	Decrease (%)
0 (control)	100	111	0
2.5	1286	339	74
5.0	1171	446	62
10.0	1130	488	58

15 **Example 2. Effects of 6,6'-dithiodinicotinic acid on mutagenesis by aminofluorene**

Number of Colonies Per Plate

2-Aminofluorene (μ g/plate)	No Inhibitor	With Inhibitor	Decrease (%)
0 (control)	149	141	0
5.0	773	705	10
15.0	2295	655	72
25.0	2134	791	63

Example 3. Effects of 6,6'-dithiodinicotinic acid on mutagenesis by 2-aflatoxinNumber of Colonies Per Plate

2-Aflatoxin (μ g/plate)	No Inhibitor	With Inhibitor	Decrease (%)
0 (control)	108	104	0
20	504	188	63
50	1117	358	68
100	1662	471	72

Example 4. Inhibition of adprp by 6,6'-dithiodinicotinic acid

ADPRP was prepared from pig thymus using the method of Khan and Shall (Biochem. Soc. Transactions 4:778 (1976)) with various modifications. Enzyme activity was approximately linear up to 15-20 minutes with NAD substrate concentration of 25 μ M.

- 10 Typical maximum enzyme activity was around 0.003 nmol product formed/minute in the assay (i.e. approximately 0.1% of substrate utilized). K_M was approximately 25 μ M.

- Using a standard assay mix containing 2mM dithiothreitol (DTT) and 100 μ M crude enzyme/ml incubation, three experiments were performed with 6,6'-dithiodinicotinic acid added at concentrations varying from 5 μ g/ml to 600 μ g/ml (i.e. 1/60th molar equivalent of DTT to 2 molar equivalents). Significant and specific inhibition of ADPRP by 6,6'-dithiodinicotinic acid was seen at 50 μ g/ml with almost complete inhibition at 150 μ g/ml and above (i.e. 0.5 molar equivalents of DTT and above). No inhibition was seen at 5 and 25 μ g/ml. The addition of excess DTT to an incubation containing 150 μ g/ml 6,6'-dithiodinicotinic acid did not increase the extent of conversion of substrate to bound product.

- 7 -

INHIBITION % OF CONTROLS

Concn of 6,6'- dithiodinicotinic acid (μ g/ml)	Percent Inhibition at:		
	7.5 min.	15 min.	20 min.
5	0	14	-
25	0	0	-
50	37	38	-
100	68	54	-
150	88	70	90
300	-	-	93
600	-	-	83

These are values for individual incubations which were stopped at either 7.5, 15, or 20 minutes. These results indicate that 6,6'-dithiodinicotinic acid is an effective inhibitor of ADPRP, forming the basis of the method of the invention for treating a human subject to inhibit mutagenic effects of carcinogens by administering to a human subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid (or derivatives thereof) to inhibit activity of ADPRP in the subjects in order to inhibit the mutagenic effects of the carcinogens. Alternatively, the invention also directs itself to a method of inhibiting ADPRP in a human subject in order to inhibit the mutagenic effects of carcinogens, the method involving administering to a human subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid (or derivatives thereof) to specifically inhibit activity of ADPRP in the subjects in order to inhibit the mutagenic effects of the carcinogens.

15

Having thus described exemplary embodiments of the present invention, it should be noted by those skilled in the art that the disclosures herein are exemplary only and that various other alternatives, adaptations, and modifications may be made within the scope of the present invention. Accordingly, the present invention is not limited to the specific embodiments as illustrated herein, but is only limited by the following claims.

20

What is claimed is:

1. A method of treating a human subject to inhibit mutagenic effects of carcinogens in said subject wherein the method comprises administering to said subject a composition
5 comprising a sufficient amount of 6,6'-dithiodinicotinic acid to specifically inhibit activity of poly(ADP-ribose) polymerase in said subject to thereby inhibit said effects.
2. The method of claim 1 wherein a sufficient amount of 6,6'-dithiodinicotinic acid is at least 0.1 mg/kilo of body weight.
- 10 3. A method of specifically inhibiting poly(ADP-ribose) polymerase in a human subject to inhibit the mutagenic effects of carcinogens on said subject, said method comprising administering to said subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid to said human to thereby inhibit said effects.
- 15 4. The method of claim 3 wherein a sufficient amount of 6,6'-dithiodinicotinic acid is at least 0.1 mg/kilo of body weight.
5. A method of inhibiting occurrence of mutations and of cancers derived from
20 mutations, which method comprises administration to a subject of a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid.
6. The method of claim 5 wherein a sufficient amount of 6,6'-dithiodinicotinic acid is at least 0.1 mg/kilo of body weight.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16262

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/44

US CL :514/335

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/335

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

HCAPLUS, USPATFULL- 6,6'-dithionicotinic acid as an antimutagenic agent and for the treatment of cancers and etc.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	Database HCAPLUS on STN, American Chemical Society, AN 1972:94870, DE 2120995 A Reversible blocking of peripheric cellular mercapto groups by 6,6'-dithiodinicotinic acid. GRASSETTI, DAVIDE R., abstract.	5-6 ----- 1-4

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

02 JANUARY 1999

Date of mailing of the international search report

20 JAN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

D. Lawrence For
M. MORZIE

Telephone No. (703) 308-1235

Alternatively, the invention also directs itself to a method of inhibiting ADPRP in a human subject in order to inhibit the mutagenic effects of carcinogens, the method involving administering to a human subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid (or derivatives thereof) to specifically inhibit activity of ADPRP in the subjects in order to inhibit the mutagenic effects of the carcinogens.

Having thus described exemplary embodiments of the present invention, it should be noted by those skilled in the art that the disclosures herein are exemplary only and that various other alternatives, adaptations, and modifications may be made within the scope of the present invention. Accordingly, the present invention is not limited to the specific embodiments as illustrated herein, but is only limited by the following claims.

What is claimed is:

1. A method of treating a human subject to inhibit mutagenic effects of carcinogens in said subject wherein the method comprises administering to said subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid to specifically inhibit activity of poly(ADP-ribose) polymerase in said subject to thereby inhibit said effects.
5
2. The method of claim 1 wherein a sufficient amount of 6,6'-dithiodinicotinic acid is at least 0.1 mg/kilo of body weight.
- 10 3. A method of specifically inhibiting poly(ADP-ribose) polymerase in a human subject to inhibit the mutagenic effects of carcinogens on said subject, said method comprising administering to said subject a composition comprising a sufficient amount of 6,6'-dithiodinicotinic acid to said human to thereby inhibit said effects.
- 15 4. The method of claim 3 wherein a sufficient amount of 6,6'-dithiodinicotinic acid is at least 0.1 mg/kilo of body weight.
5. A method of inhibiting occurrence of mutations and of cancers derived from mutations, which method comprises administration to a subject of a composition
20 comprising a sufficient amount of 6,6'-dithiodinicotinic acid.
6. The method of claim 5 wherein a sufficient amount of 6,6'-dithiodinicotinic acid is at least 0.1 mg/kilo of body weight.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/16262

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/44

US CL :514/335

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/335

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

HCAPLUS, USPATFULL- 6,6'-dithionicotinic acid as an antimutagenic agent and for the treatment of cancers and etc.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database HCAPLUS on STN, American Chemical Society, AN 1972:94870, DE 2120995 A Reversible blocking of peripheral mercapto groups by 6,6'-dithiodinicotinic acid. GRASSETTI, DAVIDE R., abstract.	5-6
-		-----
Y		1-4

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

02 JANUARY 1999

Date of mailing of the international search report

20 JAN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

M. MOEZIE

Telephone No. (703) 308-1235